



TITLE:

P-wave λn - ΣN coupling and the spin-orbit splitting of $\Lambda^9\text{Be}$

AUTHOR(S):

Fujiwara, Yoshikazu; Kohno, M.; Suzuki, Y.

CITATION:

Fujiwara, Yoshikazu ...[et al]. P-wave λn - ΣN coupling and the spin-orbit splitting of $\Lambda^9\text{Be}$. Modern Physics Letters A 2009, 24(11-13): 1031-1034

ISSUE DATE:

2009

URL:

<http://hdl.handle.net/2433/84860>

RIGHT:

c 2009 World Scientific Publishing Company.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

Modern Physics Letters A
© World Scientific Publishing Company

P-WAVE ΛN - ΣN COUPLING AND THE SPIN-ORBIT SPLITTING OF ${}^9_{\Lambda}\text{Be}$

Y. FUJIWARA¹, M. KOHNO², Y. SUZUKI³

¹ *Department of physics, Kyoto University, Kyoto 606-8502, Japan*
yfujiwar@scphys.kyoto-u.ac.jp

² *Physics Division, Kyushu Dental College, Kitakyushu 803-8580, Japan*

³ *Department of Physics, and Graduate School of Science and Technology, Niigata University*
Niigata 950-2181, Japan

Received (Day Month Year)

Revised (Day Month Year)

We reexamine the spin-orbit splitting of ${}^9_{\Lambda}\text{Be}$ excited states in terms of the SU_6 quark-model baryon-baryon interaction. The previous folding procedure to generate the $\Lambda\alpha$ spin-orbit potential from the quark-model ΛN LS resonating-group kernel predicted three to five times larger values for $\Delta E_{\ell s} = E_x(3/2^+) - E_x(5/2^+)$ in the model FSS and fss2. This time, we calculate $\Lambda\alpha$ LS Born kernel, starting from the LS components of the nuclear-matter G -matrix for the Λ hyperon. This framework makes it possible to take full account of an important P -wave ΛN - ΣN coupling through the antisymmetric $LS^{(-)}$ force involved in the Fermi-Breit interaction. We find that the experimental value, $\Delta E_{\ell s}^{\text{exp}} = 43 \pm 5$ keV, is reproduced by the quark-model G -matrix LS interaction with a Fermi-momentum around $k_F = 1.0 \text{ fm}^{-1}$, when the model FSS is used in the energy-independent renormalized RGM formalism. On the other hand, the model fss2 gives too large splitting of almost 200 keV, owing to the uncanceled contribution of the scalar-meson exchange LS components.

Keywords: Quark-model baryon-baryon interaction; spin-orbit splitting of Λ hypernuclei

PACS Nos.: 21.45.-v, 13.75.Ev, 21.80.+a, 12.39.Jh

1. Introduction

In view of rich experimental data accumulated for the light Λ -hypernuclei,^{1,2} it is important to examine if various models of the fundamental hyperon-nucleon (YN) interactions can reproduce these experimental data or not. For few-body systems, this program is most reliably carried out by detailed Faddeev calculations for the hypertriton (${}^3_{\Lambda}\text{H}$),³ ${}^4_{\Lambda}\text{H}$ and ${}^4_{\Lambda}\text{He}$,⁴ using some versions of the Nijmegen models⁵ and Jülich potentials.⁶ The knowledge of the ΛN interaction learned from these calculations, however, is mainly about the central part of the interaction and features of the ΛN - ΣN coupling of the ${}^3S_1 + {}^3D_1$ state due to the one-pion exchange tensor force. For the p -shell Λ -hypernuclei, some kinds of models inevitably need to be assumed so far, to connect properties of the Λ -hypernuclei and the under-

2 *Y. Fujiwara, M. Kohno, Y. Suzuki*

lying YN interactions. For example, the small spin-orbit (ℓs) splitting commonly observed in many of the light Λ -hypernuclei² is typically manifested in the excited states of ${}^9_\Lambda\text{Be}$, for which a simple $\Lambda + \alpha + \alpha$ three-cluster model is usually employed with appropriate $\Lambda\alpha$ and $\alpha\alpha$ potentials.⁷ In the framework of this model, the origin of the ℓs splitting for the $5/2^+$ and $3/2^+$ excited states is the spin-orbit potential between Λ and one of the α clusters, which is known to be very small due to the strong cancellation between the symmetric (LS) and antisymmetric ($LS^{(-)}$) LS forces of the ΛN interaction.

In our previous study of the ${}^9_\Lambda\text{Be}$ spectrum,⁸ we have carried out the $\Lambda\alpha\alpha$ three-cluster Faddeev calculation, trying to reproduce the very small ℓs splitting of the $5/2^+$ and $3/2^+$ excited states, $\Delta E_{\ell s}^{\text{exp}} = 43 \pm 5$ keV,² experimentally observed. As a first step, Ref. 8 directly used the quark-model (QM) ΛN LS resonating-group kernel (RGM kernel) to generate the $\Lambda\alpha$ LS potential by a simple procedure of the α -cluster folding. In this approach, the QM ΛN LS interaction of FSS or fss2 predicts 3 to 5 times larger values for $\Delta E_{\ell s}$, which is not much improved in comparison with the results of Nijmegen simulated potentials.⁷ It was pointed out in Ref. 8 that a further reduction is possible in the model FSS, if one can properly take into account the short-range correlation of the P -wave ΛN - ΣN coupling by the $LS^{(-)}$ force. This was conjectured through the analysis of the Scheerbaum factors for the single-particle (s.p.) spin-orbit potentials, calculated in the G -matrix formalism.

2. Calculational Procedure

Following the above suggestion, we here generate $\Lambda\alpha$ LS Born kernel from the LS component of the nuclear-matter G -matrix for the Λ hyperon. Our calculation consists of the following three steps.

1. Solve the G -matrix equation for the Λ -hyperon in symmetric nuclear matter with an appropriate Fermi momentum k_F and determine the s.p. potentials for N , Λ and Σ .^{9,10}
2. The LS components of the ΛN G -matrices with definite momenta K and starting energies ω are converted to the $\Lambda\alpha$ Born kernel by the folding procedure recently developed for the $\Lambda\alpha$ system.¹¹
3. Solve $\Lambda\alpha\alpha$ three-cluster system in the Faddeev formalism for composite particles.⁸

We generate $\Lambda\alpha$ LS Born kernel from our QM baryon-baryon interactions, FSS and fss2.¹² For the $(0s)^4$ α -cluster folding, a new method developed in Ref. 11 is used to derive the direct and knock-on terms of the interaction Born kernel from the ΛN G -matrix, with explicit treatments of the nonlocality and the center-of-mass motion between Λ and α . The G -matrix calculations are carried out by assuming a constant value of the Fermi momentum, $k_F = 1.07, 1.20$, and 1.35 fm⁻¹ (the normal saturation density ρ_0), since the local density approximation does not seem to work in light nuclear systems. The G -matrix equation is solved for the energy-

P-wave ΛN - ΣN Coupling and the Spin-Orbit Splitting of ${}^9_\Lambda\text{Be}$ 3

Table 1. The Scheerbaum factor S_Λ for symmetric nuclear matter and the ℓs splitting of the ${}^9_\Lambda\text{Be}$ excited states predicted by the quark-model G -matrix $\Lambda\alpha$ LS Born kernel. In the last column, “ ΛN Born” implies the previous results,⁸ in which the ΛN single-channel RGM kernel is used for the S_Λ calculation and the α -cluster folding.

	ρ/ρ_0 k_F (fm $^{-1}$)	0.5 1.07	0.7 1.20	1 1.35	ΛN Born –
G -matrix	fss2 (cont)	–11.8	–12.1	–12.3	–10.9
S_Λ (MeV fm 5)	FSS (cont)	–4.1	–5.2	–6.3	–7.8
Faddeev	fss2 (cont)	206	216	223	198
$\Delta E_{\ell s}$ (keV)	FSS (cont)	55	85	114	137
$\Delta E_{\ell s}^{\text{exp}}$ (keV)		43 ± 5			

independent QM baryon-baryon interaction, by using the renormalized RGM kernel,¹³ and the continuous prescription for intermediate spectra. A similar procedure of the renormalized RGM is also used for the microscopic $\alpha\alpha$ interaction,¹⁴ for which the Pauli forbidden states between the two α -clusters are completely eliminated in the three-cluster RGM formalism using the two-cluster RGM kernels.

3. Results and Discussion

Table 1 shows the ℓs splitting of the ${}^9_\Lambda\text{Be}$ excited states, predicted by the $\Lambda\alpha\alpha$ Faddeev calculations, using the QM G -matrix $\Lambda\alpha$ LS Born kernel. The Scheerbaum factor S_Λ is also listed to indicate the strength of the spin-orbit potentials of the Λ hyperon in symmetric nuclear matter. The Fermi momenta $k_F = 1.07, 1.20$, and 1.35 fm $^{-1}$ correspond to the densities $\rho = 0.5\rho_0, 0.7\rho_0$, and ρ_0 , respectively, with $\rho_0 = 0.17$ fm $^{-3}$ being the normal saturation density. The final values for the ℓs splitting of the $5/2^+$ and $3/2^+$ excited states are $\Delta E_{\ell s} = 55 - 114$ keV for FSS and $206 - 223$ keV for fss2, depending on the k_F values in the range of $1.07 - 1.35$ fm $^{-1}$. A smaller k_F value gives a smaller ℓs splitting. If we compare these results with the experimental value $\Delta E_{\ell s}^{\text{exp}} = 43 \pm 5$ keV, we find that the model FSS can reproduce the experimental value if the k_F value around 1.02 fm $^{-1}$ is used. We find the strong cancellation between the LS and $LS^{(-)}$ forces taking place in the QM Fermi-Breit interaction by the P -wave ΛN - ΣN coupling in the 1P_1 - 3P_1 state, when the G -matrix equation is solved especially in low-density nuclear matter. This is most prominently exhibited in the model FSS. The spin-orbit contribution from the effective-meson exchange potentials in fss2 does not lead to the sufficiently small ℓs splitting of the Λ hyperon, since the scalar-meson exchange LS force contains only the ordinary LS and does not produce the $LS^{(-)}$ force.

4. Summary

We have carried out $\Lambda\alpha\alpha$ Faddeev calculations by employing the $\Lambda\alpha$ LS Born kernel generated from the LS components of the nuclear-matter G -matrix for the Λ

4 Y. Fujiwara, M. Kohno, Y. Suzuki

hyperon. One of our SU_6 QM baryon-baryon interactions, FSS, can reproduce the very small ℓs splitting of ${}^9_\Lambda\text{Be}$ excited states, $\Delta E_{\ell s}^{\text{exp}} = 43 \pm 5$, when an appropriate k_F value corresponding to almost half of the normal saturation density is employed in the G -matrix calculation. The explicit value of k_F depends on the model construction even within the framework of the $\Lambda\alpha\alpha$ cluster model; $k_F = 1.02 \text{ fm}^{-1}$ for the model FSS, when the energy-independent renormalized RGM kernels are used for the $\alpha\alpha$ RGM kernel and for the QM baryon-baryon interaction. On the other hand, the model fss2 gives too large splitting of almost 200 keV, which is traced back to the uncanceled contribution of the scalar-meson exchange LS components. An essential ingredient of the present formalism is to take into account an important P -wave ΛN - ΣN coupling through the antisymmetric $LS^{(-)}$ force involved in the Fermi-Breit interaction. The present results indicate that the spin-orbit contribution from the effective meson-exchange potentials in fss2 needs to be improved to reproduce the small spin-orbit interaction of the Λ hyperon in the nuclear medium. A new model for the ΛN interaction with consistent central and LS components is strongly desired.

Acknowledgments

This work was supported by Grants-in-Aids for Scientific Research (C) (Grant Nos. 18540261 and 17540263), and for Scientific Research on Priority Areas (Grant No. 20028003), and Bilateral Joint Research Projects (2006-2008) from the Japan Society for the Promotion of Science (JSPS). This work was also supported by the Grant-in-Aid for the Global COE Program “The Next Generation of Physics, Spun from Universality and Emergence” from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan.

References

1. H. Bandō, T. Motoba, and J. Žofka, *Int. J. of Mod. Phys. A* **5**, 4021 (1990).
2. O. Hashimoto and H. Tamura, *Prog. Part. Nucl. Phys.* **57**, 564 (2006).
3. K. Miyagawa and W. Glöckle, *Phys. Rev. C* **48**, 2576 (1993).
4. A. Nogga, H. Kamada, and W. Glöckle, *Phys. Rev. Lett.* **88**, 172501 (2002).
5. Th. A. Rijken and Y. Yamamoto, *Phys. Rev. C* **73**, 044008 (2006).
6. J. Haidenbauer and Ulf-G. Meißner, *Phys. Rev. C* **72**, 044005 (2005).
7. E. Hiyama, M. Kamimura, T. Motoba, T. Yamada, and Y. Yamamoto, *Phys. Rev. Lett.* **85**, 270 (2000).
8. Y. Fujiwara, M. Kohno, K. Miyagawa, and Y. Suzuki, *Phys. Rev. C* **70**, 047002 (2004).
9. M. Kohno, Y. Fujiwara, T. Fujita, C. Nakamoto, and Y. Suzuki, *Nucl. Phys.* **A674**, 229 (2000).
10. Y. Fujiwara, M. Kohno, T. Fujita, C. Nakamoto, and Y. Suzuki, *Nucl. Phys.* **A674**, 493 (2000).
11. Y. Fujiwara, M. Kohno, and Y. Suzuki, *Nucl. Phys.* **A784**, 161 (2007).
12. Y. Fujiwara, Y. Suzuki, and C. Nakamoto, *Prog. Part. Nucl. Phys.* **58**, 439 (2007).
13. Y. Fujiwara, Y. Suzuki, M. Kohno, and K. Miyagawa, *Phys. Rev. C* **77**, 027001 (2008).
14. Y. Suzuki, H. Matsumura, M. Orabi, Y. Fujiwara, P. Descouvemont, M. Theeten, and D. Baye, *Phys. Lett.* **B659**, 160 (2008).